

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : G01N 24/08, 33/543	A1	(11) International Publication Number: WO 91/17428 (43) International Publication Date: 14 November 1991 (14.11.91)
(21) International Application Number: PCT/US91/03072 (22) International Filing Date: 3 May 1991 (03.05.91) (30) Priority data: 518,567 3 May 1990 (03.05.90) US (71) Applicant: ADVANCED MAGNETICS INC. [US/US]; 61 Mooney Street, Cambridge, MA 02138-1038 (US). (72) Inventors: JOSEPHSON, Lee ; 14 Churchill Avenue, Arlington, MA 02174 (US). MENZ, Edward, T. ; 17 Carle Road, Quincy, MA 02170 (US). GROMAN, Ernest, V. ; 80 Columbia Street, Brookline, MA 02146 (US). (74) Agents: SUNSTEIN, Bruce, D. et al.; Bromberg & Sunstein, 10 West Street, Boston, MA 02111 (US).		(81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SOLVANT MEDIATED RELAXATION ASSAY SYSTEM (57) Abstract A new ligand binding assay is based upon measurements of relaxation rates of the solvent, which are obtained with a magnetic resonance (MR) spectrometer. It is termed a solvent mediated relaxation assay system (SMRAS). SMRAS is based on the observation that the enhancement of proton relaxation rates produced by a magnetic material can be modulated by the binding of various analytes to a magnetic material. Hence the relaxation rate of the solvent can be interpreted to give information on the concentration of an analyte.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

SOLVENT MEDIATED RELAXATION ASSAY SYSTEM

Technical Field

5 The present invention relates to assays using magnetic materials in the presence of a magnetic field and electromagnetic radiation.

Background Art

10 Magnetic materials have been utilized in binding assays in a number of ways.

A. Magnetic Particles as Vehicle to Separate Free and Bound Labels

One of first applications of magnetic materials to immunoassays involved the attachment of antibodies to
15 magnetic particles (United States patents 4,554,088, for an invention of Whitehead; 4,141,687, for an invention of Forrest; 4,661,408, for an invention of Lau; and 3,933,997, for an invention of Hersh). This permits separation of the magnetic antibody from bulk solution, by the application of
20 an external magnetic field. After this separation has been accomplished, the amount of material attached to the magnetic antibody is determined. Quantification of the amount bound is accomplished by the use of a label or tracer attached to an antigen. Tracers of varying types have been
25 used, including radioisotopes, enzymes, chemiluminescent labels, fluorescent groups, etc. In this type of application magnetic particles serve as a vehicle for separation. Assays requiring the separation of bound and free labels, followed by quantification of the amount of
30 label bound, are termed heterogenous immunoassays or heterogenous ligand binding assays. In this type of application the magnetic property conferred on the antibody by attachment to a magnetic particle serves to facilitate separation. The same reagents could be used in an assay
35 without a magnet; for example centrifugal force can replace magnetic force as a means of separating magnetic particles from solution.

- 2 -

The magnetic particles for previously described heterogeneous immunoassays are of about a micron in size or larger. The requirement for magnetic materials of at least this size results from the fact that the force on a magnetic particle exerted by an applied magnetic field and magnetic field gradient is proportional to the mass of the magnetic material per particle. Therefore very small particles experience very weak forces from an applied magnetic field. Very small materials move too slowly to be used in assays requiring separation.

B. Magnetic Particles as Labels In Heterogeneous Immunoassay Systems.

A second use of magnetic materials in immunoassays assays involves its use as a label in a heterogeneous type of ligand binding assay (United States Patent 4,219,335, for an invention of Ebersole). In example 1 of this patent an antibody to human IgG is immobilized to a nylon film. The immobilized antibody is exposed to analyte (human IgG), which binds the immobilized antibody. After separation and washing of the film to remove unreacted analyte, anti-IgG tagged with magnetic particles is added. After the magnetic anti-IgG has been allowed to bind to the immobilized analyte (IgG), separation and washing removes unreacted, unbound magnetic particles. The result is a tertiary complex (that forms if analyte is present and the magnetic particle on the nylon film is detected in a suitable instrument). This tertiary complex may be described as follows:

30 Immobilized antibody-human IgG-magnetic antiIgG

where the symbol (-) denotes a noncovalent, potentially reversible association between two entities.

35 In its use of separation of free and bound analyte, the protocol of Ebersole is an example of a heterogeneous immunoassay. A magnetic antibody is used as a label and is

- 3 -

detected on the solid phase after separation of bound and free ligand. The magnetic particle acts as a label, replacing radioactive, enzymatic, chemiluminescent or fluorescent labels.

5 C. Electron Spin Resonance (ESR) Immunoassay

ESR immunoassay involves the covalent attachment of materials which include an unpaired electron to drugs or hormones. The resulting compound is referred to as a spin-labeled drug or spin-labeled hormone. Most often these are
10 free radicals, and most commonly a cyclical nitroxide group. Often the analyte detected in ESR immunoassay is a drug. When an antibody to a drug binds the spin-labelled drug derivative, the movement of the spin label in solution is altered, and this altered motion results in changes in the
15 ESR spectrum. Competition of the drug (analyte) prevents binding of the spin-labeled drug derivative, in the usual manner of competitive immunoassays. Since no separation is required, ESR immunoassay is a homogeneous type of assay. Electron spin immunoassay has recently been reviewed, see
20 Krick, L.J. "Ligand Binder Assays, Labels and Analytic Strategies" Marcel Dekker, New York, 1985 pp. 233-239.

Principles and Summary of the Invention

The present invention provides a ligand binding assay based upon measurements of relaxation rates of the solvent,
25 which are obtained with a magnetic resonance (MR) spectrometer. It is termed a solvent mediated relaxation assay system (SMRAS). SMRAS is based on the observation that the enhancement of proton relaxation rates produced by a magnetic material can be modulated by the binding of
30 various analytes to a magnetic material. Hence the relaxation rate of the solvent (i.e., of the protons in water, for example) can be interpreted to give information on the concentration of an analyte.

It has been often observed that the addition of
35 magnetic materials having a positive magnetic susceptibility to a solvent increases the relaxation rate(s) of nuclei measured with a magnetic resonance (MR) spectrometer. [See

- 4 -

G.L. Wolf et al. in Magnetic Resonance Annual 1985, H.Y. Kressel, ed. Raven Press, NY, 1985, pp. 231-266.] The ability to increase relaxation rates is the physical property of magnetic materials which makes them useful as contrast agents in MR imaging. Instruments such as the MR spectrometer measure the relaxation rates of nuclei in a sample and are used to assess the ability of magnetic materials to alter relaxation rates. Water is the most general solvent and the effect of a magnetic material on the relaxation of water protons is most often measured. The relaxation of protons is of two types, the so-called spin-lattice relaxation rate denoted $1/T_1$, or the spin-spin relaxation, denoted $1/T_2$. In SMRAS the binding of an analyte to a magnetic material alters the effect the magnetic material exerts on solvent relaxation rates, which are measured and interpreted to determine the concentration of analyte.

Many different kinds of biological and nonbiological molecules can be utilized in SMRAS. SMRAS often utilizes a molecule of biological origin and modifies the molecule through the attachment of a magnetic moiety, capable of altering solvent relaxation. Examples of biological molecules that can be utilized in this manner include carbohydrates, lectins, antibodies, receptors, antigens, proteins, binding proteins, nucleic acids, drugs, pesticides, etc. The magnetic moiety utilized in SMRAS can be any material capable of altering solvent relaxation as discussed below. Examples of nonbiological molecules that can be utilized in SMRAS include metal ion chelators.

In many cases, the changes in solvent relaxation rates that are measured in SMRAS reflects the formation of (or the dissolution of) large, extended complexes consisting of many magnetic particles and many analyte molecules. (The term "particle" is used to mean any material above about 5 nm in diameter.) Such complexes form in example 3, where addition of the analyte dextran breaks up the complex between Con A, a multivalent lectin binding dextran, and a dextran-coat d

- 5 -

superparamagnetic iron oxide. Similarly in example 4, bovine serum albumin (BSA) acts to dissolve or break up a complex between antibody to BSA and a BSA coated superparamagnetic iron oxide.

5 Some versions of SMRAS are similar to conventional immunoassays that are based on the formation or dissolution of large, extended macromolecular complexes. Complexes between antigens and antibodies are used in nephelometry; the complexes between antibodies and red blood cells are
10 used in hemagglutination assays; complexes between antibody coated latex particles and antigens are used in particle counting immunoassays. SMRAS is different from such conventional assays in that it uses the magnetic resonance properties of the solvent to detect complex formation, and
15 does not use radiation at or near the frequency of visible or ultraviolet light.

In some cases, the binding reactions of SMRAS do not involve the formation of large macromolecular aggregates. In example 1 magnesium (the analyte) replaces manganese in
20 complexation with a chelator, without formation of a macromolecular complex. In all cases SMRAS uses the change in solvent relaxation rates exerted by a magnetic material to determine the concentration of an analyte.

In SMRAS, an MR spectrometer is used to measure the
25 relaxation rates of water protons. For a review of MR and ESR spectrometers see Willard, H.H. et al. "Instrumental Methods of Analysis" Van Nostrand, Princeton, 1965, pp. 160-195. An MR spectrometer places the sample in a uniform applied magnetic field, and subjects it to pulses of
30 radiofrequency radiation, applied at a specific frequency, referred to as the Larmor frequency of the water protons. The signal obtained from the water protons resulting from absorption of the pulse of radiation is recorded, and the loss of signal (decay) is analyzed to obtain the
35 spin-lattice relaxation rate, and/or the spin-spin relaxation rate.

- 6 -

The distinguishing feature of SMRAS is that analyte concentration is determined from the effects a magnetic material exerts on the relaxation rates of solvent molecules. Hence we term our invention a solvent mediated relaxation assay system. In SMRAS the electromagnetic radiation interacts with the solvent, rather than with a complex of any sort between the analyte and a molecule that binds the analyte. Thus, in many conventional ligand binding assays, the incident radiation interacts directly with a complex formed between the analyte and an antibody or binding protein. Conventional ligand binding assays of this type include nephelometry, hemagglutination, latex agglutination assays and others.

The frequency of the electromagnetic radiation used in SMRAS, however, is matched to the Larmor precession frequency of the solvent, which in turn is determined by the magnetic field of the MR spectrometer. Because the radiation is at the Larmor precession frequency of the solvent, and not matched to the properties of the magnetic label employed, SMRAS can utilize many different types of magnetic materials. Though water is the most common solvent and can be used in SMRAS, other solvents might be employed containing either protons or other MR active nuclei. In an alternative embodiment, a small molecule might be added to the water whose relaxation would be enhanced by the magnetic binding element. For example, ions including the nuclei of certain isotopes (e.g. $^{23}\text{Na}^+$), phosphorous (e.g. $\text{H}_2^{31}\text{PO}_4^-$) and fluorine (e.g. $^{19}\text{F}^-$) exhibit nuclear magnetic resonance and have a high isotopic abundance. All are compatible with aqueous solvents. (Of course, other ions may be used, provided that they include the indicated isotopes of these elements or other isotopes of suitable elements exhibiting nuclear magnetic resonance.) The relaxation rates of such nuclei, like the relaxation rates of solvent protons, are enhanced by magnetic materials exhibiting a positive magnetic susceptibility. For each of the above nuclei, the incident radiation used must be at the appropriate Larmor

- 7 -

frequency, so it may be absorbed. In the case of this small-molecule mediated relaxation assay, as in the case of the solvent mediated assay, the assay is not dependent on the interaction of incident radiation with a complex between the analyte and a magnetic material. For example, in example 4 below, incident radiation does not react with the resultant magnetic BSA-antibody complex. In each case, the assay depends on the change in relative rate of the nuclei in the solvent or in the small-molecule included in the solvent exerted by the magnetic material.

The electromagnetic radiation used in MR applications, and in SMRAS, is of very low frequency (1 to 100 MHz). Radiation at this frequency is extremely safe, as the widespread use of clinical magnetic resonance imaging indicates. It should not be confused with radiation at other frequencies (10^{11} to 10^{12} MHz), that pose safety concerns.

A principal difference between SMRAS and ESR assay methods lies in the frequency of the radiation used. The ESR spectrum is obtained when the sample is placed in a magnetic field and subjected to electromagnetic radiation in the microwave range (10^4 - 10^6 MHz). The radiation used in SMRAS is of far lower frequency (1 - 10^2 MHz) than employed by ESR spin immunoassay. The frequency of radiation used in ESR relative to MR based SMRAS is related to a fundamental physical fact: the difference between energy states of unpaired electrons is far greater than the difference between energy states for nuclei exhibiting magnetic resonance.

In contrast to ESR immunoassay techniques, SMRAS uses radiation that is absorbed by the protons of the nonmagnetic (actually diamagnetic) solvent. In ESR immunoassay techniques, the radiation used is absorbed by the unpaired electron of the spin-labeled drug, which may or may not be bound to the antibody, depending on the concentration of analyte present.

- 8 -

Unlike prior art using magnetic separation strategies, the magnetic materials of the current invention are not separated from solution by an external magnetic field, during the assay. In fact many of the magnetic materials
5 used in the invention, including paramagnetic ions, paramagnetic chelates, paramagnetic proteins and superparamagnetic colloids (below the micron size range), cannot be separated from solution in a timely fashion with the magnetic fields produced by common permanent magnets.

10 Brief Description of the Drawings

The foregoing features of the invention may be more readily understood by reference to the following detailed description taken with the accompanying drawings, in which;

Figure 1 plots concentration of the analyte (magnesium)
15 in example 1 below versus MR relaxation time of the solvent; and

Figure 2 plots concentration of the analyte (glucose) in example 2 below versus MR relaxation time of the solvent;

Figure 3 plots concentration of the analyte (dextran)
20 in example 3 below versus MR relaxation time of the solvent;

Figure 4 plots concentration of the analyte (BSA) in example 4 below versus MR relaxation time of the solvent;

Figure 5 plots concentration of the analyte (DTPA) in example 5 below versus MR relaxation time of the solvent.

25 Figure 6 shows the effect of inorganic phosphate on the $1/T_2$ produced by a 5 mM solution of $MnCl_2$.

Figure 7 shows the decrease in $1/T_2$ relaxation rate and increase in absorbance with time that results from the activity of alkaline phosphatase.

30 Detailed Description of Specific Embodiments

A. Utility of SMRAS

SMRAS has major advantages over other types of ligand binding assays.

First, SMRAS can work on opaque fluids, whole blood,
35 milk, lipemic plasma, hemolyzed blood, etc. With conventional ligand assays, the optical quality of the samples necessitates preparation to remove interfering

- 9 -

substances, before the ligand binding assay protocol can commence.

To obtain the relaxation rates of a solvent, SMRAS utilizes an MR spectrometer. This instrument places the sample in a strong magnetic field and subjects it to radiofrequency radiation at the so-called Larmor frequency of the solvent protons. SMRAS utilizes electromagnetic radiation in the radiofrequency range rather than in the visible range, for the detection of the ligand binding.

The indifference of SMRAS to properties of the solvent is due to the fact that the radiofrequency radiation used in the measurement of proton relaxation penetrates most biological systems. In MR imaging the penetrability of radiofrequency radiation is used to advantage. For example the protons in bone marrow proteins can be excited with radiofrequency radiation that has passed through bone and an MR image of bone marrow obtained.

Second, SMRAS is a homogeneous type assay. That is, the assay protocol does not involve a separation of bound and free ligands. This lack of a separation step results in a more user friendly, less labor intensive assay procedure.

Third, SMRAS can measure widely different types of materials as analytes, including materials with different molecular weights and different physicochemical properties.

In example 1, below, an assay for a low molecular weight divalent cation (magnesium), is demonstrated. In example 2 an assay for a monosaccharide, in this case glucose, is demonstrated. In example 3 an assay for a polysaccharide polymer, in this case dextran, is demonstrated. In example 4 an assay for protein of about 65,000 daltons (BSA) is demonstrated.

Fourth, a wide variety of magnetic materials including superparamagnetic iron oxide colloids, paramagnetic chelates, paramagnetic ions, paramagnetic red blood cells, or ferromagnetic particles, can be used in SMRAS. These vastly different materials all exhibit a positive magnetic susceptibility, and because of their positive magnetic

- 10 -

susceptibility increase relaxation rates of solvent protons. By altering their effect on solvent relaxation rates, all can be utilized in SMRAS. The ability of SMRAS to utilize different types of magnetic materials is shown in the

5 examples below. In example 1, paramagnetic manganese ion, which has a molecular weight of less than 100 daltons, is used to alter solvent relaxation. In example 3 a dextran coated superparamagnetic iron oxide colloid, which has a molecular weight in excess of 500,000 daltons, is used. The

10 superparamagnetic iron oxide possesses a far higher magnetic susceptibility than the paramagnetic manganese ion, although both may be employed.

Fifth, SMRAS offers the possibility of performing multiple ligand binding assays simultaneously, by utilizing

15 many of the data reduction and instrumentation techniques developed for MR imaging. For example, in MR imaging the magnitude of the applied magnetic field can be varied, so that only the protons in different samples resonate at slightly different Larmor frequencies. In MR imaging this

20 is referred to as slice selective excitation. In SMRAS the corresponding technique involves simultaneously placing several sample tubes in a spectrometer that applies a slightly different magnetic field to each sample. As a result protons in each sample will resonate at different

25 Larmor frequencies. The relaxation rates of multiple samples can be determined simultaneously.

There are several general strategies for synthesizing magnetic materials for use in the present invention.

1. Coupling paramagnetic ions to drugs, proteins,

30 carbohydrates, nucleic acids, or other biologically active groups:

Paramagnetic ions are often attached to molecules by derivatization of the molecule with a chelator, followed by allowing the chelate to bind a paramagnetic ion. Examples

35 of this approach include Mears et al. Anal. Biochem. 142:68-78 (1984); Mears and Wensel, Acc. Chem. Res.

- 11 -

17:202-209 (1984); Khaw t al. Science 209:295-297; Ogan et al, Invest. Radiol. 22:665-671 (1987).

2. Coupling magnetic particles or magnetic colloids to drugs, proteins, carbohydrates, nucleic acids, or
5 biologically active groups:

There are a variety of techniques for attaching molecules to magnetic particles and/or colloids such as United States Patents 4,141,687, for an invention of Forrest; 4,661,408, for an invention of Lau; 4,452,773, for
10 an invention of Molday; 4,795,698, for an invention of Owen; 4,554,088, for an invention of Whitehead; and 3,993,997, for an invention of Hersh.

In the present invention examples 2, 3 and 4 utilize magnetic colloids.

- 15 3. Using red blood cells as magnetic "particles":

The paramagnetism exhibited by the deoxyhemoglobin of red blood cells permits the cells to act as magnetic particles, i.e. red blood cells filled with deoxyhemoglobin alter solvent relaxation rates obtained with the MR
20 technique (Thulborn et al. Biochim. Biophys. Acta 714:265-270 (1982)). Assays of red blood cell surface antigens are the basis of blood typing performed prior to blood transfusion. Such assays often involve the agglutination of red blood cells by antibodies, with an
25 optical measurement of agglutination. This red blood cell magnetic agglutination is analogous to the aggregation or dispersion of other magnetic labels used in SMRAS. Thus red blood cells can be utilized in a SMRAS assay without chemical modification to confer a positive magnetic
30 susceptibility upon them.

4. Measuring the rate of an enzyme reaction:

A valuable feature of SMRAS demonstrated by the examples above is its ability to measure the concentration of analytes by utilizing assays based on ligand binding. In
35 ligand binding assays antibodies, lectins or other binding proteins bind the analyte and the degree of binding is used in the estimation of analyte concentration. However, SMRAS

- 12 -

can also measure enzymatic reaction rates, even though the molecules that serve as substrates (or which are produced as products) of enzymatic reactions are usually diamagnetic and have no significant effect on the relaxation rate of a solvent. For example, the common enzyme alkaline phosphatase hydrolyzes a substrate paranitrophenol phosphate (PNP) into the products, paranitrophenol and inorganic phosphate (PO_4^{3-} or Pi). Neither substrate nor products are paramagnetic or exert significant effects on solvent relaxation. However, one of the products of the reaction, Pi, forms a complex with manganese ion (Mn^{2+}), which greatly increases solvent relaxation rates over that characteristic of pure solvent alone. Thus, Pi reacts with Mn^{2+} to form a complex which reduces the $1/T_2$ of the solvent produced by Mn^{2+} .

Figure 6 shows the effect of increasing concentrations of Pi as NaH_2PO_4 on a solution of 5 mM MnCl_2 . At approximately equimolar concentrations of Pi and Mn^{2+} , the $1/T_2$ decreases to 5 sec^{-1} , which is about that of water. In other words, the addition of Pi eliminates the effect the Mn^{2+} has on $1/T_2$, i.e. the effect on solvent relaxation.

Figure 7 shows an enzymatic assay for alkaline phosphatase based on the principle described above. After addition of alkaline phosphatase to a solution of 5 mM MnCl_2 and 10 mM PNP, paranitrophenol and Pi are formed. The accumulation of paranitrophenol is monitored by its absorbance at 410 nm. The accumulation of Pi is evident from its reaction with Mn^{2+} and by the fall in $1/T_2$. Assays for alkaline phosphatase in serum are widely used in clinical chemistry, and alkaline phosphatase has been attached to antibodies and used as a general label in immunoassays.

In the example above, the common substrate PNP is used in conjunction with MnCl_2 , to provide an enzyme assay according to the principles of SMRAS. Thus, no special synthetic chemistry was used to demonstrate the applicability of SMRAS to the assay of alkaline phosphatase.

- 13 -


An alternative approach would be to design and synthesize molecules as substrates for SMRAS, much the way the chromogenic substrates are synthesized for spectrophotometric or fluorometric assay systems. For example, a suitable substrate could be one which when converted into a product(s) has very different effects on solvent relaxation.

B. Examples

Example 1: Assay for Magnesium Ion.

10 Background and Principle of the Method:

Paramagnetic Mn^{2+} is a powerful enhancer of proton relaxation. When manganese is bound to a chelating compound such as diethylenetriaminepentaacetic acid, DTPA, its ability to enhance proton relaxation is greatly reduced. Magnesium ion (Mg^{2+}) is a diamagnetic ion that also binds to DTPA, but which has virtually no effect on proton relaxation, whether bound to DTPA or not. When added to appropriate concentrations of DTPA and manganese, magnesium prevents the binding of manganese to DTPA as shown below.



As a result the concentration of manganese ion is increased and an increase in $1/T_2$ results.

Assay Procedure:

To a test tube was added 0.5 mL of 25 mM DTPA adjusted to pH 8.5 with NaOH, and 0.5 mL of 12.5 mM $MnCl_2$. $MgCl_2$ at various concentrations (0.2 mL) was then added. The solutions were vortexed and the value of $1/T_2$ determined. The standard curve of Figure 1 was obtained. All relaxation rate measurements are obtained in an IBM PC-20 MR spectrometer at 39°C, using the Carr Purcell Meiboom Gill (CPMG) technique.

Example 2: Assay for Glucose.

Background and Principle of the Method:


Concanavalin A (Con A) is a lectin that binds carb hydrates such as dextran. The dextran coated superparamagnetic iron oxide was synthesized by the method of Molday, see above. Monosaccharides like glucose can

- 14 -

bind Con A and prevent Con A from binding a dextran coated superparamagnetic iron oxide.

When con A binds a dextran coated superparamagnetic iron oxide, the effect the colloid exerts on $1/T_2$ increases at low concentrations of Con A but then decreases at higher concentrations (data not shown). In this example concentrations of Con A and superparamagnetic iron oxide have been chosen to maximize the effect the colloid exerts on $1/T_2$. Under these conditions the addition of glucose breaks up the Con A-dextran coated superparamagnetic iron oxide complex, and reduces $1/T_2$. This is represented below. (The symbol (/) indicates a permanent bond between two materials under conditions of the assay, while the symbol (-) indicates a dissociable bond.)

15



20 Assay Procedure:

Half a mL of Con A (1 mg/mL) and 0.5 mL of dextran coated superparamagnetic iron oxide (0.8 mM iron) were mixed together in a test tube and then 0.2 mL glucose, 0 to 200 mg/mL, was added. After 20 minutes at 37 °C, relaxation rates were measured as in example 1. The response of the assay to glucose is shown in Figure 2.

Example 3: Assay for Dextran.

Background and Principle of the Method:

The procedure is as in Example 2, except that the dextran replaces glucose in favoring the dissociation of the complex between dextran coated superparamagnetic iron oxide and Con A.

Assay Procedure:


The procedure of example 2 is followed except that a dextran T-10, replaces glucose. The response of the assay to dextran is shown in Figure 3.

- 15 -

Example 4: Assay For Bovine Serum Albumin (BSA) Using Antibody to BSA.

Background and Principle of the Method:

A BSA coated superparamagnetic iron oxide colloid is
5 utilized and reacted with antibody to BSA. Concentrations
of antibody and colloid are chosen such that the presence of
antibody decreases the effect the colloid exerts on $1/T_2$.
Addition of BSA favors the dissociation of the antibody from
the BSA coated superparamagnetic iron oxide. The addition
10 of BSA increases the value of $1/T_2$. The reaction might be
represented:



15 Assay Procedure:

A solution of BSA (0.1 mL) at the indicated
concentration in phosphate buffered saline with 0.15%
gelatin (PBS-gel) is added to a test tube. Next 0.1 mL of
BSA coated superparamagnetic iron oxide colloid is added.
20 The colloid used was the BSA-coated particle from Immunicon
Corp., (Huntington Valley PA). Finally, 0.1 mL of anti-BSA
is added. The antibody was diluted 1:40 in PBS-gel and was
from Sigma Chemical (St. Louis, MO).

The mixture is incubated at 37 °C for 60 minutes and
25 the relaxation rate ($1/T_2$) of the 300 mL sample determined
as above.

The standard curve for the assay is shown in Figure 4.

Example 5: Assay for Chelating Equivalent Groups on a Protein.


30 Background and Principle of the Method:

In this example DTPA anhydride is reacted with the
protein fetuin from calf serum, to give a high molecular
weight protein chelator denoted Fet/DTPA. The assay
involves ascertaining the capacity of the modified prot in
35 (Fet/DTPA) to decrease $1/T_2$ by comparing its effect with
solutions of DTPA in known concentrations. This is
represented below.

- 16 -



OR



5 Assay Procedure:

1. Synthesis of Fet-DTPA

Fifty mg of fetuin is dissolved in 10 mL of 0.1M NaHCO₃, pH 8 buffer and 120 mg of DTPA anhydride added. Unreacted DTPA was removed by dialysis against the same buffer using a
10 dialysis bag with a 10 kilodalton cutoff.

2. Assay Procedure:

To a test tube is added 0.2 mL of 1 mM MnCl₂, 0.2 mL DTPA solutions at known concentrations (0- 5600 nM) or 0.2 mL of Fet-DTPA (2mg/mL in fetuin but with unknown amount of
15 DTPA). The reaction mixture is incubated at room temperature for 30 minutes after which values of 1/T₂ are determined. The standard curve shown in Figure 5 was obtained for DTPA. The number of equivalent chelating groups on the protein was determined from the effect of a
20 comparable concentration of DTPA.

- 17 -

What is claimed is:

1. A method, of assaying the concentration of an analyte in a solvent, comprising:

using a magnetic material having positive magnetic
5 susceptibility to alter the relaxation characteristics of a nucleus included in the solvent, such relaxation characteristics obtained by a magnetic resonance technique, the magnetic material being selected so that its effect, on the relaxation of the nucleus included in the solvent, is
10 modified by the binding of the analyte to the magnetic material; and

measuring the relaxation characteristics of the nucleus by a magnetic resonance technique.

2. A method according to claim 1, wherein the
15 magnetic material is a magnetic metal oxide colloid or magnetic particle, with a polymer-coated surface.

3. A method according to claim 1, wherein the magnetic material is a protein with a paramagnetic ion attached.

20 4. A method according to claim 1, wherein the magnetic material is a red blood cell.

5. A method according to claim 1, wherein the magnetic material is a paramagnetic ion.

6. A method according to claim 1, wherein the
25 magnetic material is a magnetic antibody.

7. A method according to claim 1, wherein the magnetic material is a magnetic antigen.

8. A method according to claim 1, wherein the nucleus included in the solvent is part of the solvent itself.

30 9. A method according to claim 2, wherein the nucleus included in the solvent is part of the solvent itself.

10. A method according to claim 3, wherein the nucleus included in the solvent is part of the solvent itself.

11. A method according to claim 4, wherein the nucleus
35 included in the solvent is part of the solvent itself.

12. A method according to claim 5, wherein the nucleus included in the solvent is part of the solvent itself.

- 18 -

13. A method according to claim 6, wherein the nucleus included in the solvent is part of the solvent itself.

14. A method according to claim 7, wherein the nucleus included in the solvent is part of the solvent itself.

5 15. A method according to claim 1, wherein the nucleus included in the solvent is an ion that is an isotope of sodium having an atomic weight of 23.

16. A method according to claim 1, wherein the nucleus included in the solvent is an ion that is an isotope of
10 phosphorous having an atomic weight of 31.

17. A method according to claim 1, wherein the nucleus included in the solvent is an ion that is an isotope of fluorine having an atomic weight of 19.

18. A method according to claim 1, wherein the analyte
15 is selected from the group consisting of a substrate of an enzyme and a product of a reaction of the enzyme.

19. A method according to claim 1, wherein the analyte is an enzyme.

20. A method according to claim 19, wherein the enzyme
20 is used as a tracer in an immunoassay to measure the concentration of an antibody or an antigen.

21. A method according to claim 1, wherein the analyte is an antigen.

22. A method according to claim 1, wherein the analyte
25 is an antibody.

23. A method, of measuring the rate of an enzyme reaction in a solvent, comprising:


assaying the concentration of a component that is a substrate or product of the enzyme reaction, by using a
30 magnetic material having positive magnetic susceptibility to alter the relaxation characteristics of a nucleus included in the solvent, such relaxation characteristics obtained by a magnetic resonance technique, the magnetic material being selected so that its effect, on the relaxation of the
35 nucleus included in the solvent, is modified by the binding of the component to the magnetic material; and

- 19 -


measuring the relaxation characteristics of the nucleus
by a magnetic resonance technique.

24. A method according to claim 23, wherein the
nucleus included in the solvent is part of the solvent
5 itself.

1/7



SUBSTITUTE SHEET



SUBSTITUTE SHEET

3/7





FIG. 3

4/7



5/7





FIG. 5

SUBSTITUTE SHEET






FIG. 7

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 91/03072

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl.5 G 01 N 24/08 G 01 N 33/543		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl.5	G 01 N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ^o	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	WO,A,8601112 (FOX CHASE CANCER CENTER) 27 February 1986, see page 4, lines 2-17; page 6, lines 14-20 ---	1-24
Y	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 254, no. 6, 25 March 1979, (US), A. RIGO et al.: "Nuclear magnetic relaxation of 19F as a novel assay method of superoxide dismutase", pages 1759-1760, see the whole article ---	1-24
A	BIOCHIMICA ET BIOPHYSICA ACTA, vol. 714, no. 2, 2 February 1982, Elsevier Biomedical Press, K.R. THULBORN et al.: "Oxygenation dependence of the transverse relaxation time of water protons in whole blood at high field", pages 265-270 (cited in the application) --- -/-	
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>^o Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
02-09-1991	01 OCT 1991 	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	MISS T. TAZELAAR	

III. DOCUMENTS CONSIDERED TO BE RELEVANT

(CONTINUED FROM THE SECOND SHEET)

Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	<p>MAGNETIC RESONANCE IN MEDICINE, vol. 3, 1986, Academic Press, Inc., C.F.G.C. GERALDES et al.: "Magnetic field dependence of solvent proton relaxation rates induced by Gd³⁺ and Mn²⁺ complexes of various polyaza macrocyclic ligands: implications for NMR imaging" -----</p>	

US 9103072
SA 47914

64M41 5NHC) 3 (0.154

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82